

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

# **Host-race specificity in the endemic pygmy mistletoe *Korthalsella* *salicornioides* (Viscaceae)**

A thesis presented in partial fulfilment of the requirements for the degree of

Master of Science

In

Plant Biology

at Massey University, Palmerston North,

New Zealand

**Sofie Margaret Pearson**

2016



## ABSTRACT

*Korthalsella* Tiegh. is a genus of stem hemiparasites in the family Viscaceae, represented in New Zealand by three endemic species: *K. clavata*, *K. lindsayi*, and *K. salicornioides*. The most host-specific is *K. salicornioides* as it parasitizes two main host genera *Leptospermum* (Myrtaceae) and *Kunzea* (Myrtaceae), while the other two species are considered generalists parasitizing a wider range of host species. *K. salicornioides* is naturally uncommon and sparse, although it can be locally abundant on occasion. Mistletoe populations are at risk primarily due to habitat destruction and subsequent loss of hosts. Cross-infection experiments in *K. salicornioides* provided some insight into the presence of putative host races, as better mistletoe seedling establishment success rates were apparent when the maternal and recipient hosts were the same. However, because previous molecular sequence data (nuclear internal transcribed spacers and chloroplast *trnQ-rps16*) for *K. salicornioides* were not informative about specific host-races, more rapidly evolving molecular markers might be expected to detect host races.

In this study, next generation sequencing was used to develop novel microsatellite markers for *Korthalsella*. Eleven markers were reliably amplifiable and the most polymorphic for *K. salicornioides* were used to genotype 272 *K. salicornioides* individuals from 16 populations. Across all populations few alleles were identified, and within-population assessment of genetic variation indicated that many populations have low levels of genetic diversity and high proportions of homozygotes. Despite the presence of few alleles, a high degree of genetic differentiation between most populations was detected and was found to reflect host species and geography.

The findings of this study that *Korthalsella salicornioides* populations have low levels of genetic variation but host-specific races, has important conservation implications. The main conservational focus should be maintaining and increasing host *Leptospermum* and *Kunzea*

populations. The spread of mistletoe seed on hosts within or between populations may also increase the chances of continued survival. However, it is imperative that genetic material comes from the same host species, and consideration should also be given to the geographic area, especially in the Wairarapa. This study provides insights into the population structure within and between the different host populations and suggests several interesting areas of future study.

## ACKNOWLEDGEMENTS

First and foremost I would like to give a humongous thanks to my supervisors, Jennifer Tate and Vaughan Symonds, for your guidance and advice along the way. Thank you for giving me the opportunity to undertake this study. The time put into helping me collect samples, answer my questions, give feedback, and notice my numerous spelling mistakes is hugely appreciated.

I would also like to express my gratitude for the help of Alastair Robertson, Amir Sultan, Nick Singers and Peter de Lange for information about *Korthalsella* and identifying populations to collect. Also to Paul Cashmore (DOC) and Conny Flanigan (Kawerau District Council) for taking time out of your day to show and help me collect from the mistletoe populations.

Thanks to all the LoST lab members, both past and present, who helped me with my study and the advice you gave me. I would like to express my gratitude to Megan van Etten for all her work and help getting me started, and to Prashant Joshi for all the help provided over the years.

I would also like to thank the Auckland Botanical Society for the Lucy Cranwell Grant, the Heseltine Trust for the Heseltine Trust Bursary and Massey University for the J. P. Skipworth Scholarship for aid in this research.

I am also grateful to my close friends and family for all their help, enthusiasm and encouragement throughout my studies. I couldn't have achieved this without everyone's support.

## ABBREVIATIONS

%P	-percentage of polymorphic loci
A	-number of alleles
A <sub>S</sub>	-allele size
A <sub>T</sub>	-total number of alleles
AFLP	-Amplified Fragment Length Polymorphism
AMOVA	-analysis of molecular variance
BLAST	-basic local alignment search tool
bp	-base pairs
cpDNA	-chloroplast DNA
CASS	-cheaply amplified size standard
CTAB	-hexa-decetylammonium bromide
DNA	-deoxyribonucleic acid
dNTP	-deoxyribonucleotide triphosphate
DOC	-Department of Conservation (New Zealand)
ER	-Ecological Region
F <sub>IS</sub>	-component of Wright's (1921) fixation index, used to define within population structure by calculating the average observed heterozygosity of an individual relative to the expected heterozygosity of individuals in the population it belongs to
F <sub>ST</sub>	-component of Wright's (1921) fixation index, used to define between population structure by comparing the expected heterozygosity of individuals within a subpopulation to the total expected heterozygosity of individuals across all populations
H <sub>E</sub>	-expected heterozygosity
H <sub>O</sub>	-observed heterozygosity
IBD	-Isolation by Distance
ITS	-internal transcribed spacer
LnP(D)	-mean posterior probability

MA	-Million years
MPN	-Dame Ella Campbell Herbarium
mtDNA	-mitochondrial DNA
N	-number of individuals
$N_A$	-number of alleles
$N_E$	-effective number of alleles
PCR	-Polymerase Chain Reaction
RAPD	-Randomly Amplified Polymorphic DNA
SNP	-Single Nucleotide Polymorphism
$T_m$	-melting temperature
VNTR	-Variable Number of Tandem Repeat loci



# TABLE OF CONTENTS

Abstract .....	iii
Acknowledgements .....	v
Abbreviations .....	vi
List of Figures.....	xi
List of Tables.....	xii
 <b>1. Introduction .....</b>	 <b>1</b>
1.1 Parasitic Flowering Plants .....	1
1.2 Mistletoes .....	5
1.3 <i>Korthalsella</i> in New Zealand .....	6
1.3.1 Conservation of <i>Korthalsella</i> and its hosts .....	10
1.3.2 Dispersal mechanisms and life history traits.....	12
1.3.3 Evidence for host-race.....	13
1.4 Markers to Assess Genetic Variation .....	17
1.5 Analysis Methods for Population Structure.....	19
1.6 Focus of this Research.....	21
1.7 References .....	22
 <b>2. Microsatellite Markers for <i>Korthalsella</i> (Viscaceae) in New Zealand.....</b>	 <b>29</b>
2.1 Abstract.....	29
2.2 Introduction .....	30
2.3 Methods and Results .....	30
2.4 Conclusions .....	34
2.5 References .....	39

<b>3. Population genetics and host-race specificity in <i>Korthalsella salicornioides</i> (Viscaceae) ...</b>	<b>41</b>
3.1 Abstract .....	41
3.2 Introduction .....	42
3.3 Materials and Methods.....	46
3.3.1 Sample collection .....	46
3.3.2 DNA extraction and genotyping .....	50
3.3.3 Assessing genetic variation.....	51
3.3.4 Resolving genetic structure and differentiation.....	52
3.4 Results.....	53
3.4.1 Genetic Variation.....	53
3.4.2 Genetic Structure .....	55
3.4.3 NeighborNet .....	62
3.5 Discussion.....	64
3.5.1 Genetic variation within <i>Korthalsella salicornioides</i> populations .....	64
<i>Private alleles</i> .....	64
<i>Seed dispersal and vector limitation of Korthalsella salicornioides</i> .....	66
<i>Lower North Island genetic “hot spot”</i> .....	68
3.5.2 Host-races in <i>Korthalsella</i> .....	70
<i>Mechanisms of parasite differentiation</i> .....	71
3.5.3 Conservation implications for <i>Korthalsella salicornioides</i> and its hosts .....	72
<i>Conservation of Korthalsella salicornioides</i> .....	73
<i>Host trees as economic benefits</i> .....	75
3.5.4 Future directions .....	77
3.6 Conclusion.....	79
3.7 References .....	81

<b>4. Conclusion.....</b>	<b>89</b>
4.1 Introduction .....	89
4.2 Findings .....	91
4.3 Limitations.....	93
4.4 Future Directions .....	94
4.5 References .....	97
 <b>Appendix 1: Population genetic diversity estimates for 16 populations of <i>Korthalsella salicornioides</i> based on the seven polymorphic loci.....</b>	 <b>98</b>

## LIST OF FIGURES

### 1. Introduction

**Figure 1.1** *Korthalsella salicornioides* parasitic on *Leptospermum scoparium* and *Kunzea tenuicaulis* ..... 8

**Figure 1.2** Primary, secondary and tertiary host of *Korthalsella salicornioides*, *K. clavata* and *K. lindsayi* ..... 9

**Figure 1.3** ITS sequence type diversity in *Korthalsella salicornioides*..... 16

### 3. Population genetics and host-race specificity in *Korthalsella salicornioides*

**Figure 3.1** Distribution map of the populations of *Korthalsella salicornioides* sampled for this study ..... 48

**Figure 3.2** Plot of  $\Delta K$  vs K for STRUCTURE results based on 15 replicates for each K value.. 58

**Figure 3.3** Plot of mean posterior probability values per cluster, based on 15 iterations per K from STRUCTURE analyses..... 58

**Figure 3.4** STRUCTURE cluster assignment of *Korthalsella salicornioides* individuals..... 59

**Figure 3.5** Mantel test results displayed in graph of pairwise  $F_{ST}(1-F_{ST})$  against the natural log of geographic distance (km) for *Korthalsella salicornioides* ..... 61

**Figure 3.6** Mantel test results displayed in graph of pairwise  $F_{ST}(1-F_{ST})$  against the natural log of geographic distance (km) for *Korthalsella salicornioides* mānuka-host populations (A), and kanuka-host populations (B)..... 61

**Figure 3.7** NeighborNet generated in SPLITSTREE4 based on pairwise genetic distances between all *Korthalsella salicornioides* populations..... 63

**Figure 3.8** *Kunzea amathicola* at Hokio Beach ..... 68

### 4. Conclusion

**Figure 4.1** Leaf shapes of herbarium specimens of *Leptospermum scoparium* parasitized by *Korthalsella salicornioides*..... 95

**Figure 4.2** Two *Leptospermum scoparium* host populations at Kohi Point coastal habitat along walking track (A), and Kerikeri swamp habitat (B) ..... 96

## LIST OF TABLES

### 1. Introduction

<b>Table 1.1</b> The 12 orders of parasitic angiosperms with example families and genera .....	3
--	---

### 2. Microsatellite Markers for *Korthalsella* (Viscaceae) Species in New Zealand

<b>Table 2.1</b> Locations and herbarium voucher information for <i>Korthalsella</i> populations used in this study .....	35
---	----

<b>Table 2.2</b> Primer sequences and characteristics of 16 microsatellite loci developed from <i>Korthalsella salicornioides</i> .....	36-37
---	-------

<b>Table 2.3</b> Genetic properties of the newly developed 11 microsatellite loci across five North Island populations of <i>Korthalsella</i> .....	38
---	----

### 3. Population genetics and host-race specificity in *Korthalsella salicornioides*

<b>Table 3.1</b> Relative <i>Korthalsella salicornioides</i> population size information based on the estimated number of host trees parasitized for locations used in this study .....	47
---	----

<b>Table 3.2</b> Locations, host tree and herbarium voucher information for <i>Korthalsella salicornioides</i> populations used in this study .....	49
---	----

<b>Table 3.3</b> Characteristics of eleven microsatellite loci for 272 samples of <i>Korthalsella salicornioides</i> .....	54
--	----

<b>Table 3.4</b> Population information and genetic diversity estimates for 16 populations of <i>Korthalsella salicornioides</i> .....	55
--	----

<b>Table 3.5</b> Clustering of K=7 STRUCTURE results examining genetic structure within the <i>Kunzea</i> and <i>Leptospermum</i> genera.....	60
---	----

<b>Table 3.6</b> AMOVA results for the partitioning of microsatellite variation in: A) all <i>Korthalsella salicornioides</i> populations, B) comparing between four regions identified in STRUCTURE .....	62
--	----